



## CLOCK GENE VARIATION IN TYPE 2 DIABETES: A REVIEW

Nushrat Parveen\*, Trilochan Satapathy, Pushpa Prasad, Amit Roy

Columbia Institute of Pharmacy, Tekari, Raipur, Chhattisgarh – 493111, India

Diabetes mellitus type 2 is a long-standing metabolic disorder that is exemplified by high blood sugar, insulin resistance, and comparative lack of insulin. General symptoms include increased thirst, frequent urination, and unsolved weight loss. Type 2 diabetes is mainly due to obesity and not sufficient work out in public who are heritably prone. Circadian clocks are significant to keep the moment in the sequence of physiological practice, series of behaviour and metabolism. The plasma level of glucose and numerous hormones implicated in glucose homeostasis for example insulin and glucagon exhibit circadian variation. Circadian desynchrony, a feature of alter occupation elevated-fat diet feed and sleep distraction in individual have been linked with metabolic disorders for instance obesity and type 2 diabetes. Circadian rhythm distraction can cause different fitness disarray. Current reading has discovered a seal connection among the pathophysiology of metabolic condition, which is characterized by obesity and hyperglycemia, and the operation of interior molecular clocks.

**Keywords:** Clock genes, Diabetes mellitus, Pathophysiology, Single nucleotide polymorphisms, Obesity

### INTRODUCTION

Every mammalian cells almost show an arrangement of genes, called as clock genes. By way of negative and positive auto-regulatory feedback loops of transcription and translation these control the circadian rhythm of cellular actions.<sup>[1,2]</sup> Clock (Circadian Locomotor Output Cycles Kaput) is a gene programming a general helix-loop-helix-PAS transcription factor (CLOCK) that change both the resolution and period of circadian rhythms. CLOCK utilizes as a vital activator of downstream components in the passageway significant to the production of circadian rhythms.<sup>[3]</sup> The uncontrollably advance in the episode of type 2 diabetes (T2D) has fetch out the term for to study in supplementary seriousness and extent the hazardous matter that put in to this sickness in array to have a supplementary wide-ranging revelation of the course to clarify the trail towards more defined and efficient precautionary attachment.<sup>[4-6]</sup> A relationship among a polymorphism close to the circadian clock gene CRY2 and increased fasting glucose have been discovered by current genome-wide connection learnings.<sup>[1,2,19]</sup>

### PRINCIPLE

Circadian clocks undergo the principle of consecutive gene commencement in the type of a rotation: the preliminary commencement of a gene is synchronized through the final one in the series building up an auto-regulatory reaction loop on behalf of which single cycle acquire on 24 h.<sup>[7]</sup>

### For Correspondence

nushratparveen77@gmail.com


### MECHANISM

In mammals the circadian clock system is through two interconnecting dogmatic reaction loops. In the foremost loop, two transcriptional activators►Bmal1 (brain and muscle ARNT-like protein 1) and► Clock (or Npas2 in neuronal tissue) figure heterodimers within the cytoplasm and make a way into the nucleus where they unite to► E-box series in the supporter of ►Period (Per1,2) and►Cryptochrome (Cry1,2) genes causal to the establishment of their appearance. In the cytoplasm a variety of array of Per and Cry proteins assist with each other, walk off into the nucleus and hinder the action of Bmal1/Clock or Bmal1/Npas2 complexes. Devoid of these complexes triggering transcription of the Per and Cry genes, level of Per and Cry transcripts and their relevant protein yield turn down therefore Per and Cry genes close up their individual transcription.<sup>[7,8]</sup> A second loop controls the appearance of the Bmal1 gene. Inside the nucleus Bmal1/Clock or Bmal1/Npas2 heterodimers unite to E-boxes present in the promoters of genes that instruct the retinoic acid-related orphan nuclear receptors► Rev-erba and► Rora, which battle for the ROR constituent (RORC) in the Bmal1 supporter. Rora stimulate Bmal1 expression, whereas Rev-erba stifle it. As an outcome oscillations of Bmal1 and Rora/Rev-erba are absent. If establishment succeed above term Bmal1 protein is formed and it outline heterodimers inside the cytoplasm among Clock or Npas2 depending on the tissue. These heterodimers penetrate the nucleus and commence the subsequent cycle of gene establishment of mutual loops.<sup>[7-9, 20]</sup>

## LINK BETWEEN CIRCADIAN CLOCKS AND METABOLISM

Circadian clocks control every day alterations within a widespread series of physiological and behavioral utility with the timing of sleep. Distraction of circadian biology or sleep has reflective metabolic consequences. There is clear association among circadian clocks and metabolism.<sup>[11]</sup> In murine models, transgenic distractions of key clock-linked genes provoke distinct metabolic phenotypes.<sup>[18]</sup> In individual never-ending desynchrony of interior circadian time through the exterior surroundings (e.g., during shift work) interlinks with elevated frequency of obesity, type 2 diabetes, and cardiometabolic disorders. The mammalian circadian timing arrangement is made of a “master” clock inside the suprachiasmatic nuclei (SCN) of the hypothalamus and a cycle of “peripheral” clocks that are there in extra-SCN province of the brain in addition to typically every additional tissues. The ecological light–dark phase harmonize the circadian organization via performing throughout the SCN clock, which after that retains temporal array of the marginal clocks through numerous periodic yield. These SCN-driven passageway involve neuronal (e.g., sympathetic character), endocrine (e.g., melatonin and cortisol emission), and behavioral (e.g., nourishing) action. Murine white adipose tissue (WAT), like by several additional tissues, hold endogenous circadian timing property and ~10–20% of murine WAT transcriptome is detected to show a 24-h deviation. It is noteworthy that mice that are heritably prone to stoutness (obesity) and type 2 diabetes or fed an elevated stout diet show decreased amplitude rhythm of gene appearance in WAT, guiding to the supposition that the sturdiness of WAT clocks is functionally linked with metabolic phenotype.<sup>[11,12]</sup> The liver plays a major task in preserving metabolic events to normal feeding–fasting cycles. This role is displayed via the circadian phrase of the entire liver genes integrated within the biotransformation of lipids, proteins, carbohydrates, and xenobiotics.<sup>[13]</sup> The pathophysiology of T2DM includes insulin resistance, pancreatic b-cell dysfunction and primitive adiposity and is compound. It has been acknowledged for decades that a distraction of biological rhythms (which occurs almost certainly amid shift work) elevates the harm of rising obesity and T2DM. The hypothalamus regulates a huge arrangement of physiological procedures, with the involvement of sleep/wake cycles, sexual deeds and reproduction and metabolic control for instance thermoregulation, energy ingestion/ outflow, glucose

metabolism, lipid metabolism, and food and water intake. All through the action/nourishing period blood glucose is principally of dietetic source and throughout the sleeping/hunger period glucose is consecutively recruited from endogenous glucose construction inside the liver to retain blood glucose intensity in a comparatively slight array. In this procedure liver glycogen substance go through huge day by day fluctuations to keep up blood glucose rank since glycogen production and deprivation be specially recruited all through the action/feeding and inactive/hunger episodes correspondingly.<sup>[14-17]</sup>



[7]




Fig. 2: Hypothetical clock mechanism in mammals<sup>[10]</sup>

## CONCLUSION

Quick steps have been made in current years to scrutinize the compound relationship among circadian rhythms, metabolism,

and sleep. These three methods are mutually dependent and intimately linked all the way through wide mutual association. Researchers have revealed over the precedent few years that cellular (or regional) clocks can be found in the liver, pancreas and different parts of the body as well. Evenly eating or sleeping at the wrong times may put off these peripheral clocks out of synchronization with the master clock in the brain. Surplus advances will emerge by refining the neuroanatomic map associating circadian and energetic centers with areas regulating sleep and wakefulness; such efforts may ultimately lead to the development of rational therapeutics for the treatment of obesity and sleep-related metabolic diseases.

## REFERENCES

1. Stamenkovic et al., "Clock genes in human islets", *Metabolism: Clinical and Experimental*, 2012, vol. 61:7, pp. 978-985
2. Marcheva B, Ramsey KM, Affinati A, Bass J (2009) Clock genes and metabolic disease. *J Appl Physiol* 107: 1638-1646
3. <https://en.wikipedia.org/wiki/CLOCK>
4. Corella et al. *Cardiovascular Diabetol* (2016) 15:4, DOI 10.1186/s12933-015-0327
5. Leon BM, Maddox TM. Diabetes and cardiovascular disease: epidemiology, biological mechanisms, treatment recommendations and future research. *World J Diabetes*. 2015;6:1246–58.
6. Pandey A, Chawla S, Guchhait P. Type-2 diabetes: current understanding and future perspectives. *IUBMB Life*. 2015;67:506–13.
7. Comp.by: Sgunasekar Date:22/3/08 Time:12:33:54 Stage:First Proof File Path://spiina1001z/womat/production/PRODEV/000000005/0000006643/0000000016/0000773876. D Proof by: QC by: Clock Genes Au1, U<sub>RS</sub> ALBRECHT, JURGEN A. RIPPERGER Department of Medicine, Division of Biochemistry, University of Fribourg, Fribourg, Switzerland
8. Young MW, Kay S (2001) Time zones: a comparative genetics of circadian clocks. *Nat Rev Genet* 2:702–715
9. Ko CH, Takahashi JS (2006) Molecular components of the mammalian circadian clock. *Hum Mol Genet* 15(2):R271–R277
10. [https://www.google.co.in/search?q=hypothetical+clock+m+echanism+in+mammals+-+images&ie=utf-8&oe=utf-8&aq=t&rls=org.mozilla:en-US:official&client=firefox-beta&channel=fflb&gfe\\_rd=cr&ei=uKDAVuORC8eM8Qf726LwB](https://www.google.co.in/search?q=hypothetical+clock+m+echanism+in+mammals+-+images&ie=utf-8&oe=utf-8&aq=t&rls=org.mozilla:en-US:official&client=firefox-beta&channel=fflb&gfe_rd=cr&ei=uKDAVuORC8eM8Qf726LwB)
11. Rhythmic Diurnal Gene Expression in Human Adipose Tissue From Individuals Who Are Lean, Overweight, and Type 2 Diabetic Daniella T. Otway, Simone Mäntele, Silvia Bretschneider, John Wright, Paul Trayhurn, Debra J. Skene, M. Denise Robertson, and Jonathan D. Johnston
12. Oishi K, Atsumi G, Sugiyama S, et al. Disrupted fat absorption attenuates obesity induced by a high-fat diet in Clock mutant mice. *FEBS Lett* 2006;580:127–130
13. David Gatfield and Ueli Schibler\*, Department of Molecular Biology and National Center for Competence in Research Frontiers in Genetics, Sciences III, University of Geneva, 30, Quai Ernest Ansermet, CH-1211 Geneva-4, Switzerland
14. Circadian control of glucose metabolism, Andries Kalsbeek, Susanne la Fleur Eric FliersÓ 2014 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY NC-ND license
15. Damiola, F., Le Minh, N., Preitner, N., Kornmann, B., Fleury-Olela, F., Schibler, U., 2000. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes & Development* 14:2950-2961
16. Kennaway, D.J., Owens, J.A., Voultsios, A., Boden, M.J., Varcoe, T.J., 2007. Metabolic homeostasis in mice with disrupted Clock gene expression in peripheral tissues. *American Journal of Physiology* 293:R1528-R1537
17. Clock Genes and Energy Metabolism H.-K. Hong et al. P.J. Shiromani et al. (eds.), *Sleep Loss and Obesity: Intersecting Epidemics* DOI 10.1007/978-1-4614-3492-4\_2, © Springer Science+Business Media, LLC 2012
18. Otway DT, Frost G, Johnston JD: Circadian rhythmicity in murine pre-adipocyte and adipocyte cells. *Chronobiol Int* 26:1340-1354, 2009
19. Lyssenko V, Nagorny CL, Erdos MR, et al. (2009) Common variant in MTNR1B associated with increased risk of type 2 diabetes and impaired early insulin secretion. *Nat Genet* 41: 82-88
20. Ko CH, Takahashi JS (2006) Molecular components of the mammalian circadian clock. *Hum Mol Genet* 15(2): R271–R277

Received 14<sup>th</sup> November 2014

Revised 21<sup>st</sup> November 2014

Accepted 26<sup>th</sup> November 2014

J. App. Pharm. Res., 3 (1); 2015: 01 – 03